

# Practice Parameters for Using Polysomnography to Evaluate Insomnia: An Update

An American Academy of Sleep Medicine Report

Standards of Practice Committee of the American Academy of Sleep Medicine

Michael Littner, MD;<sup>1</sup> Max Hirshkowitz, PhD;<sup>2</sup> Milton Kramer, MD;<sup>3</sup> Sheldon Kapen, MD;<sup>4</sup> W. McDowell Anderson, MD;<sup>5</sup> Dennis Bailey, DDS;<sup>6</sup> Richard B. Berry, MD;<sup>7</sup> David Davila, MD;<sup>8</sup> Stephen Johnson, MD;<sup>9</sup> Clete Kushida, MD, PhD;<sup>10</sup> Daniel I. Loube, MD;<sup>11</sup> Merrill Wise, MD;<sup>12</sup> B. Tucker Woodson, MD<sup>13</sup>

<sup>1</sup>VA Greater Los Angeles Healthcare System and UCLA School of Medicine, Sepulveda, CA; <sup>2</sup>Baylor College of Medicine and VA Medical Center, Houston, TX; <sup>3</sup>Maimoides Medical Center, Psychiatry Department, Brooklyn, NY and New York University School of Medicine, New York, NY; <sup>4</sup>VA Medical Center and Wayne State University, Detroit, MI; <sup>5</sup>College of Medicine, University of South Florida, Tampa, FL; <sup>6</sup>Englewood, Colorado; <sup>7</sup>University of Florida Health Science Center, Gainesville, FL; <sup>8</sup>Baptist Medical Center, Little Rock, AR; <sup>9</sup>St. Patrick Hospital Sleep Center, Missoula, Montana; <sup>10</sup>Stanford University Center of Excellence for Sleep Disorders, Stanford, CA; <sup>11</sup>Sleep Medicine Institute, Swedish Medical Center, Seattle, WA; <sup>12</sup>Departments of Pediatrics and Neurology, Baylor College of Medicine, Houston, TX; <sup>13</sup>Department of Otolaryngology and Communication Sciences, Medical College of Wisconsin, Milwaukee, WI.

**Abstract:** Insomnia is a common and clinically important problem. It may arise directly from a sleep-wake regulatory dysfunction and/or indirectly result from comorbid psychiatric, behavioral, medical, or neurological conditions. As an important public-health problem, insomnia requires accurate diagnosis and effective treatment. Insomnia is primarily diagnosed clinically with a detailed medical, psychiatric, and sleep history. Polysomnography is indicated when a sleep-related breathing disorder or periodic limb movement disorder is suspected, initial diagnosis is uncer-

tain, treatment fails, or precipitous arousals occur with violent or injurious behavior. However, polysomnography is not indicated for the **routine** evaluation of transient insomnia, chronic insomnia, or insomnia associated with psychiatric disorders.

**Citation:** Standards of Practice Committee of the American Academy of Sleep Medicine. Practice parameters for using polysomnography to evaluate insomnia: an update for 2002. *SLEEP* 2003;26(6):754-60.

## INTRODUCTION

INSOMNIA IS CHARACTERIZED BY A COMPLAINT OF DIFFICULTY INITIATING SLEEP, MAINTAINING SLEEP, AND/OR NONRESTORATIVE SLEEP THAT CAUSES CLINICALLY SIGNIFICANT DISTRESS OR IMPAIRMENT IN SOCIAL, OCCUPATIONAL, OR OTHER IMPORTANT AREAS OF FUNCTIONING. Insomnia is associated with a variety of morbidities including decreased quality of life, absenteeism, auto accidents, and increased general health care utilization. The term *insomnia* is used to describe a wide range of alterations in the amount and type of sleep loss or perceived sleeplessness. Etiologies include insomnia provoked directly by intrinsic sleep disorders, extrinsic sleep disorders, or circadian rhythm irregularities. Additionally, insomnia is associated with many psychiatric, medical, and neurological conditions. Insomnia may also be due to the use or misuse of either prescription or over-the-counter medications. In total, 55 of the nosological entities listed in The International Classification of Sleep Disorders<sup>1</sup> can have insomnia as a symptom. Epidemiological research reveals that insomnia afflicts 10-30 percent of the adult population.<sup>2-4</sup>

Overnight polysomnography (CPT 95810) is a standard tool in sleep medicine for evaluating sleep-related pathophysiology, sleep architecture, and sleep integrity. Some etiologies underlying insomnia have specific pathophysiology detectable with polysomnography (e.g. periodic limb movements). Other insomnias may manifest abnormal sleep architectural patterns (e.g. major depressive disorder) that while recognizable are diagnostically nonspecific.<sup>5-7</sup> Finally, sleep integrity can be directly measured with polysomnography. Measures such as latency to sleep onset, total sleep time, number of arousals and awakenings, and sleep efficiency are routinely calculated to characterize a night of sleep.<sup>8,9</sup> Disturbance in such measures objectively verify complaints of difficulty initiating and maintaining sleep. Furthermore, polysomnographic criteria can differentiate physiologically-based sleep disturbances from sleep state misperception and helps to evaluate whether the subject pos-

sibly prefers the drug for the wrong reasons (e.g. because of euphoriant properties). For this reason, polysomnography is a component of the standard procedure used to verify insomnia and assess treatment efficacy for research purposes.<sup>10</sup>

Some clinicians in their search for objective diagnostic indices have used polysomnography to evaluate patients with insomnia. Monitoring for specific etiology-related pathophysiologies (e.g. obstructive sleep apnea) can be very useful for making a diagnosis when insomnia is secondary to another condition. By contrast, using sleep integrity measures to diagnose primary insomnia has several potential drawbacks. First, most normal sleepers experience transient insomnia (or some degree of sleep disruption) the first time they sleep in the laboratory. This *first night effect* disappears when the individual acclimates to the novel sleep environment. Second, insomnia typically varies in severity across nights such that a single night may fail to properly characterize the full extent of the sleep problem.<sup>11,12</sup> Third, patients with psychophysiological insomnia often paradoxically sleep well on their first night in the laboratory.<sup>13</sup> Fourth, some patients with insomnia have sleep state misperception; that is, they have the complaint of insomnia in the absence of objective findings from polysomnography.<sup>14-17</sup> These four factors make it difficult to achieve diagnostic specificity with a single night of polysomnography. For these reasons, cost for multiple nights in the laboratory are generally considered prohibitive unless a clear advantage over other procedures is provided.

Thus, using polysomnography to diagnose primary insomnia is fraught with methodological pitfalls detracting from its usefulness. As such, using polysomnography for **routine** clinical evaluation of transient or chronic insomnia is controversial and was not recommended in the 1995 American Sleep Disorders Association Practice Parameter Report.<sup>18,19</sup> By contrast, diagnostic polysomnography was endorsed as sometimes appropriate in cases where a) diagnosis is uncertain, b) sleep-related breathing disorder or periodic limb movement disorder are suspected, c) a patient is refractory to treatment, d) violent behaviors are

comorbid, or e) circadian dysrhythmias complicate the clinical picture.

This update modifies and replaces the 1995 practice parameters.

## METHODS

The Standards of Practice Committee appointed a task force to review the role of polysomnography in the evaluation of insomnia. The task force examined the previously published practice parameters and the reviews upon which they were based. Excluding conference abstracts and letters to the editor, the references cited in the Reite et al (1995)<sup>19</sup> review paper were considered in the current literature reassessment. Medline was searched from 1980 through and including articles published up to February 2002. The terms *insomnia*, *sleeplessness*, and *sleep initiation and maintenance disorders* were crossed with the terms *polysomnography*, *sleep evaluation*, *monitoring ambulatory*, or *monitoring physiologic*. Searches were also conducted crossing *Ekbom's syndrome*, *restless legs syndrome*, *nocturnal myoclonus*, *fibromyalgia*, and *depression with polysomnography*, *sleep evaluation*, *monitoring ambulatory*, or *monitoring physiologic*. The two Medline searches were then combined and limited to human subject and English language publications. Data from the articles were extracted, reviewed, and summarized.

Based on the evidence review, updated practice recommendations were developed by the Standards of Practice Committee. Recommendations are rated as standards, guidelines, or options (Table 1) based on evidence from studies published in peer-reviewed journals that were evaluated and listed in the evidence tables (Tables 3A, 3B, and 3C). However, when scientific evidence is not available, insufficient, or inconclusive, the recommendations were based on consensus opinion of the committee.

These practice parameters define principles of practice that should meet the needs of most patients in most situations. These guidelines are neither inclusive of all proper methods of care nor exclusive of other methods of care reasonably directed toward obtaining the same result. Judgment regarding the propriety of any care strategy ultimately must be made by health care providers with consideration given to individual circumstances presented by the patient, available diagnostic procedures, and extant treatment resources.

The Board of Directors of the American Academy of Sleep Medicine approved these recommendations. All members of the American Academy of Sleep Medicine Standards of Practice Committee and Board of Directors completed detailed conflict-of-interest statements and were found to have no conflicts of interest with regard to this subject.

The American Academy of Sleep Medicine expects these guidelines to have a positive effect on professional behavior, patient outcomes, and possibly health care costs. These practice parameters reflect the state of knowledge at the time of development and will be reviewed, updated, and revised, as new information becomes available.

## RESULTS

The Medline search strategy produced 706 citation titles. Two Standards of Practice Committee (SPC) members (MH and MK) reviewed all of the titles, abstracts, and if needed the full publication to

**Table 1**—AASM Recommendations (adapted from Eddy<sup>56</sup>)

Term	Description
<i>Standard</i>	A <i>standard</i> is a generally accepted patient care strategy reflecting a high degree of clinical certainty. The term <i>standard</i> generally implies a basis in either Level I evidence directly addressing the clinical issue or overwhelming Level II evidence.
<i>Guideline</i>	A <i>guideline</i> is a patient care strategy reflecting a moderate degree of clinical certainty. The term <i>guideline</i> implies a basis in Level II evidence or a consensus of Level III evidence.
<i>Option</i>	An <i>option</i> is a patient care strategy reflecting uncertain clinical use. The term <i>option</i> implies either inconclusive or conflicting evidence, or conflicting expert opinion.

determine whether the citation was appropriate for this review. Inclusion criteria were: 1) the study included laboratory polysomnography, 2) a formal diagnosis was rendered, and 3) there appeared to be a focus on insomnia. Exclusion criteria were: 1) reports of treatment outcome studies, 2) reports of ambulatory PSG and specialized waveform analysis, 3) reports that did not present original data (editorials, letters, and reviews), and 4) single case studies. One reviewer (MK) selected 28 papers while the other (MH) selected 34 papers; there were 23 articles selected by both reviewers and these were distributed among 4 SPC members (MH, MK, SK, and MA) for data extraction. One additional article was excluded because it turned out to be a review. Two of the 4 SPC members independently extracted information from each article. Extraction discrepancies were resolved with extraction or grading by another SPC member. Only minimal discrepancies were encountered. Additionally, each article was graded according to criteria [modified from Sackett (1993)<sup>20</sup>] shown in table 2. The extracted data was then summarized in Tables 3A, 3B, and 3C. Of the 8 articles summarized in Table 3A that address diagnostic utility of polysomnography in patients with insomnia, 5 were level III, grade C; 2 were level IV, grade C, and 1 was level V, grade D. Table 3B focuses on polysomnography in psychiatric disorders associated with insomnia. Except for a single paper rated as level III, grade C, the remaining 7 articles on this topic were graded as level IV, grade C. Finally, Table 3C tabulates the 6 other relevant reports. One each was rated as level III, grade C and level V, grade D; the remaining 4 articles were level IV, grade C. It is apparent that there is little rigorous scientific work regarding diagnostic polysomnographic evaluation of insomnia. Most of the studies are un-blinded, cross-sectional, non-randomized with comparisons to historical, opportunistic, or case controls. Sample sizes for comparative or case control work are generally limited; however, a few larger sample case series have been published.

Articles were evaluated to address the utility of polysomnography for the diagnosis of insomnia according to whether there was evidence to answer the following questions. The results are detailed in Tables 3A, B and C.

1. Did polysomnography help in the evaluation of insomnia in this study?
2. Did polysomnography provide information helpful for understanding treatment failure?
3. Did polysomnography differentiate between insomnia of different etiologies?

## RECOMMENDATIONS

When possible, practice recommendations are evidence based. New recommendations, as well as those that are the same as, similar to, or expansion of, previous ones are noted in the text.

**Table 2**—AASM classification of evidence (Adapted from Sackett, 1993)<sup>20</sup>

Grades	Evidence Levels	Study Design
A	I	1-Blind, Prospective Study 2-Large sample with a spectrum of patients reviewed 3-Results are compared according to a reference standard
B	II	1-Blind, Prospective Study 2-Limited sample or limited spectrum of patients included 3-Results are compared according to a reference standard
C	III	1-Not blind, randomized, or prospective 2-Methodologically limited 3-Results are compared according to a reference standard
C	IV	1-Not blind, randomized, or prospective 2-Methodologically limited 3-Results not compared according to a reference standard
D	V	1-Not blind, randomized, or prospective 2-Methodologically limited 3-Results not compared to any reference

**1. Insomnia is an important public-health problem that requires accurate diagnosis and effective treatment. (Standard)**

Insomnia is a symptom of an underlying disorder or condition. The insomnia may be a problem directly related to the sleep-wake regulatory system and/or it might be associated with a comorbid psychiatric, behavioral, medical, or neurological condition. This recommendation is similar to the one made previously and is based on committee consensus.<sup>2,18,19,21,22</sup>

**2. Insomnia is primarily diagnosed by clinical evaluation through a careful, detailed medical, psychiatric, and thorough sleep history (which includes assessment of sleep patterns and waking processes). (Standard)**

This recommendation is similar to the one previously made and is based on committee consensus.<sup>18,19,21,23</sup> The change from previous recommendation is that there is a greater emphasis on the sleep history.

**3. Polysomnography is indicated when sleep-related breathing disorders or periodic limb movement disorder is suspected. (Standard)**

This recommendation is similar to one previously made (differing primarily in wording) and is based on 4 level III (grade C) studies,<sup>24-27</sup> 1 level IV (grade C) study,<sup>28</sup> work cited in the previous practice parameter,<sup>22,29,30</sup> and committee consensus.

**4. Polysomnography is indicated when initial diagnosis is uncertain, treatment fails (behavioral or pharmacologic), or precipitous arousals occur with violent or injurious behavior. (Guideline)**

This recommendation was made previously and is based on work cited in the previous practice parameter<sup>31-38</sup> and committee consensus.

**5. Polysomnography is not indicated for the routine evaluation of transient or chronic insomnia. (Guideline)**

This is the same recommendation as made previously<sup>18,39</sup> (with minor wording changes) and is based on committee consensus.

**6. Polysomnography is not indicated for the routine evaluation of insomnia due to psychiatric disorders. (Guideline)**

This recommendation modifies the one previously made and is based on 1 level III (grade C) study,<sup>40</sup> 7 level IV (grade C) studies<sup>41-49</sup> and committee consensus.

**7. Polysomnography is not clinically useful in differentiating the insomnia associated with dementia from other forms of insomnia, including insomnia associated with depression. (Guideline)**

This recommendation is the same as one previously made and is based on 1 level IV (grade C) study,<sup>50</sup> 1 level V (grade D) study,<sup>51</sup> work cited in the previous practice parameter<sup>34-38</sup> and committee consensus.

**8. Polysomnography is not useful in establishing the diagnosis of insomnia associated with fibromyalgia or chronic fatigue syndrome because the alpha-delta sleep pattern described in fibromyalgia syndrome is a nonspecific finding. (Guideline)**

This recommendation is the same as one previously made and is based on 1 level IV (grade C) study,<sup>52</sup> work cited in the previous practice parameter,<sup>53-55</sup> and committee consensus.

**FURTHER RESEARCH**

This evidence based review revealed significant weakness in the published literature concerning the diagnostic utility of polysomnography for clinically evaluating patients with insomnia. More methodologically sound research, with attention to diagnostic sensitivity and specificity is needed. Large scale, prospective, controlled studies are needed to assess the sensitivity and specificity of polysomnographic metrics to diagnose insomnia. Such studies must assure blind data recording, scoring, and interpretation. Computerized microarchitectural analysis, if directed specifically at improving diagnostic discrimination may be helpful in difficult cases. Furthermore, blind, prospective, controlled studies are needed to evaluate polysomnography's ability to differentiate between insomnias differing in etiologies and comorbidities. Well-delineated subgroups for specific secondary insomnias, as well as, subgroups for different insomnias *lumped* together as primary insomnia need to be considered. Issues surrounding perceived sleeplessness without objective findings, so-called *sleep state misperception*, needs particular attention. Progress has already been made in moving research away from using heterogeneous samples labeled *insomnia* and has yielded some progress. Finally, econometric analysis of diagnostic practice should be undertaken. Such research is a priority because insomnia is a serious, prevalent condition that adversely affects productivity, morbidity, mortality, and quality of life. Ultimately, improved diagnostic practice will improve the overall quality of care.

**REFERENCES**

1. ICSD- International classification of sleep disorders: Diagnostic and coding manual. Diagnostic Classification Steering Committee. Thorpy MJ, Chairman. Rochester, Minnesota: American Sleep Disorders Association, 1990.
2. Ohayon MM, Caulet M, Priest RG, et al. DSM-IV and ICSD-90 insomnia symptoms and sleep dissatisfaction. *Br J Psychiat* 1997;171:382-8.
3. Blazer DG, Hays JC, Foley DJ et al. Sleep complaints in older adults: a racial comparison. *J Gerontol A Biol Sci Med Sci* 1995; 50: M280-2
4. Foley DJ, Monjan AA, Brown SL et al. Sleep complaints among elderly persons: an epidemiologic study of three communities. *Sleep* 1995; 18: 425-32.
5. Hauri PJ. A cluster analysis of insomnia. *Sleep* 1983;6:326-38.
6. Hudson JI, Pope HG, Sullivan LE, Waternaux CM, Keck PE, Broughton RJ. Good sleep, bad sleep: a meta-analysis of polysomnographic measures in insomnia, depression, and narcolepsy. *Biol Psychiatry* 1992;32:958-75.
7. Merica H, Gaillard JM. The EEG of the sleep onset period in insomnia: a discriminant analysis. *Physiol Behav* 1992;52:199-204.
8. Standards of Practice Committee of the American Sleep Disorders Association, Practice parameters for the use of polysomnography in the evaluation of insomnia. *Sleep* 18; 55-7: 1995.
9. Rechtschaffen A, Kales A. A manual of standardized terminology, techniques and scoring system for sleep stages of human sleep subjects. Washington, DC: U.S. Government Printing Office, 1968.
10. Parrino L, Terzano MG. Polysomnographic effects of hypnotic drugs. A review. *Psychopharmacology* 1996; 126(1):1-16.
11. Coates TJ, Killen JD, George J, et al. Discriminating good sleepers from insomniacs using all-night polysomnograms conducted at home. *J Nerv Ment Disease* 1982;170:224-30.
12. Coates TJ, George JM, Killen JD, Marchini E, Hamilton S, Thorenson CE. First night effects in good sleepers and sleep-maintenance insomniacs when recorded at home. *Sleep* 1984;4:293-8.
13. Hauri PJ, Olmstead EM. Reverse first night effect in insomnia. *Sleep* 1989;12:97-105.
14. Baekeland F, Hoy P. Reported vs. recorded sleep characteristics. *Arch Gen Psychiatry* 1971; 24: 548-51.
15. Carskadon MA, Dement WC, Mitler MM, Guilleminault C, Zarcone VP, Spiegel R. Self-report versus laboratory findings in 122 drug-free subjects with complaints of chronic insomnia. *Am J Psychiatry* 1976; 33:1382-8.
16. Frankel BL, Coursey RD, Buchbinder R, Snyder F. Recorded and reported sleep in chronic primary insomnia. *Arch Gen Psychiatry* 1976;33:615-23.
17. Salin-Pascual RJ, Roehrs TA, Merloiti LA, Zorick F, Roth T. Longterm study of the sleep of insomnia patients with sleep state misperception and other insomnia patients. *Am J Psychiatry* 1992;147:904-8.
18. Sleep Disorders Atlas Task Force of the American Sleep Disorders Association. EEG arousals: scoring rules and examples. *Sleep* 1992;15:173-84.
19. Reite M, Buysse D, Reynolds C, Mendelson W. The use of polysomnography in the evaluation of insomnia. *Sleep* 1995;18:58-70.
20. Sackett D. Rules of evidence and clinical recommendation. *Can J Cardiol* 1993; 9: 487-9.
21. ASDA Diagnostic Classification Steering Committee, Thorpy MJ, chairman. International classification of sleep disorders: diagnostic and coding manual. Rochester, MN: American Sleep Disorders Association, 1990
22. Coleman RM, Roffwarg HP, Kennedy SJ, et al. Sleep-wake disorders based on a

- polysomnographic diagnosis: a national cooperative study. *J Am Med Assoc* 1982;247:997-1003.
23. Sateia M J, Doghramji K, Hauri PJ, Morin CM. Evaluation of chronic insomnia. An American Academy of Sleep Medicine review. *Sleep* 2000;23:243-308.
  24. Vgontzas AN, Kales A, Bixler EO, Manfredi RL, Vela-Bueno A. Usefulness of polysomnographic studies in the differential diagnosis of insomnia. *International J Neurosci* 1995; 82:47-60.
  25. Lichstein KL, Riedel BW, Lester KW, Aguillard RN. Occult sleep apnea in a recruited sample of older adults with insomnia. *J Consulting & Clin Psychol* 1999; 67:405-10.
  26. Roehrs T, Conway W, Wittig R, Zorick F, Sickelsteel J, Roth T. Sleep-wake complaints in patients with sleep-related respiratory disturbances. *Am Rev Respiratory Disease* 1985; 132:520-3.
  27. Zorick FJ, Roth T, Hartzel KM, Piccione PM, Stepanski EJ. Evaluation and diagnosis of persistent insomnia. *Am J Psychiat* 1981; 138:769-73.
  28. Reynolds CF 3<sup>rd</sup>, Coble PA, Black RS, Holzer B, Carroll R, Kupfer DJ. Sleep disturbances in a series of elderly patients: polysomnographic findings. *J Am Geriatrics Soc* 1980; 28:164-70.
  29. Edinger JD, Hoelscher TJ, Webb MD, Marsh GR, Radtke RA, Erwin CW. Polysomnographic assessment of DIMS: empirical evaluation of its diagnostic value. *Sleep* 1989; 12:315-22.
  30. Kales A, Bixler EO, Soldatos CR, Vela-Bueno A, Caldwell AB, Cadieux RJ. Biopsychobehavioral correlates of insomnia, part 1: role of sleep apnea and nocturnal myoclonus. *Psychosomatics* 1982; 23: 589-600.
  31. Dyken ME, Rodnitzky RL. Periodic, aperiodic and rhythmic motor diseases of sleep. *Neurology* 1992;42(7, Suppl 6):68-74.
  32. Hauri PJ, Friedman M, Ravaris CL. Sleep in patients with spontaneous panic attacks. *Sleep* 1989;12:323-37.
  33. Jacobs EA, Reynolds CF, Kupfer DJ, Lovine PA, Ehrenpreis AB. The role of polysomnography in the differential diagnosis of chronic insomnia. *Am J Psychiatry* 1988;145:346-9.
  34. Prinz PN, Reskind ER, Vitaliano PP, et al. Changes in the sleep and waking EEGs of nondemented and demented elderly subjects. *J Am Geriatr Soc* 1982;30:86-93.
  35. Prinz PN, Vitaliano PP, Vitiello MV, et al. Sleep EEG and mental function changes in senile dementia of the Alzheimer's type. *Neurobiol Aging* 1982;3:361-70.
  36. Reynolds CF, Kupfer DJ, Hoch CC, et al. Sleep deprivation as a probe in the elderly. *Arch Gen Psychiatry* 1987;44:982-90.
  37. Vitiello MV, Bokan JA, Kukull WA, Muniz RL, Smallwood RG, Prinz PN. Rapid eye movement sleep measures of Alzheimer's-type dementia patients and optimally healthy aged individuals. *Biol Psychiatry* 1984;19:721-34.
  38. Vitiello MV, Bliwise DL, Prinz PN. Sleep in Alzheimer's disease and the sundown syndrome. *Neurology* 1992;42(Suppl 6):83-94.
  39. Standards of Practice Committee of the American Academy of Sleep Medicine. Practice parameters for the evaluation of chronic insomnia. *Sleep* 2000; 23 :237-41.
  40. Thase ME, Kupfer DJ, Fasiczka AJ, Buysse DJ, Simons AD, Frank E. Identifying an abnormal electroencephalographic sleep profile to characterize major depressive disorder. *Biological Psychiatry* 1997; 41:964-73.
  41. Arriaga F, Paiva T, Matos-Pires A, Cavaglia F, Lara E, Bastos L. The sleep of non-depressed patients with panic disorder: a comparison with normal controls. *Acta Psychiatrica Scandinavica* 1996; 93:191-4.
  42. Gillin JC, Duncan WC, Pettigrew KD, Frankel BL, Snyder F. Successful separation of depressed, normal, and insomniac subjects by EEG sleep data. *Arch Gen Psychiatry* 1979;36:85-90.
  43. Haynes SN, Follingstad DR, McGowan WT. Insomnia: sleep patterns and anxiety level. *J Psychosom Res* 1974;18:69-74.
  44. Hohagen F, Lis S, Krieger S, et al. Sleep EEG of patients with obsessive-compulsive disorder. *European Arch Psychiatry & Clin Neurosci* 1994; 243:273-8.
  45. Hurwitz TD, Mahowald MW, Kuskowski M, Engdahl BE. Polysomnographic sleep is not clinically impaired in Vietnam combat veterans with chronic posttraumatic stress disorder. *Biological Psychiatry* 1998; 44:1066-73.
  46. Lauer CJ, Krieg JC, Garcia-Borreguero D, Ozdaglar A, Holsboer F. Panic disorder and major depression: a comparative electroencephalographic sleep study. *Psychiatry Research* . 1992; 44:41-54.
  47. Merica H, Blois R, Bovier P, Gaillard JM. New variables for defining sleep continuity. *Physiology & Behavior* 1993; 54:825-31.
  48. Pecknold JC, Luthe L. Sleep studies and neurochemical correlates in panic disorder and agoraphobia. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*. 1990; 14:753-8.
  49. Stein MB, Enns MW, Kryger MH. Sleep in nondepressed patients with panic disorder: II. Polysomnographic assessment of sleep architecture and sleep continuity. *J Affective Disorders* 1993; 28:1-6.
  50. Allen SR, Seiler WO, Stahelin HB, Spiegel R. Seventy-two hour polygraphic and behavioral recordings of wakefulness and sleep in a hospital geriatric unit: comparison between demented and nondemented patients. *Sleep* 1987; 10:143-59.
  51. Evans BM., Bartlett JR. Prediction of outcome in severe head injury based on recognition of sleep related activity in the polygraphic electroencephalogram. *J Neuro Neurosurgery & Psychiatry* 1995; 59:17-25.
  52. Manu P, Lane TJ, Matthews DA, Castriotta RJ, Watson RK, Abeles M. Alpha-delta sleep in patients with a chief complaint of chronic fatigue. *Southern Med J* 1994; 87:465-70.
  53. Hauri P, Hawkins DR. Alpha-delta sleep. *Electroencephalogr Clin Neurophysiol* 1973;34:233-7.
  54. Mahowald MW, Mahowald ML, Bundlie SR, Ytterberg SR. Sleep fragmentation in rheumatoid arthritis. *Arth Rheum* 1989;32:974-83.
  55. Moldofsky H, Scarisbrick P, England R, Smythe H. Musculoskeletal symptoms and non-REM sleep disturbance in patients with "fibrositis syndrome" and healthy subjects. *Psychosom Med* 1975;37:341-51.
  56. Reprinted with permission from American College of Physicians. Eddy DM. *A Manual for Assessing Health Practices and Designing Practice Policies: The Explicit Approach*. Philadelphia: American College of Physicians, 1992.
  57. Dorsey CM, Bootzin RR. Subjective and psychophysiological insomnia: an examination of sleep tendency and personality. *Biological Psychiatry* 1997; 41:209-16.
  58. Ferini-Strambi L, Filippi M, Martinelli V, et al. Nocturnal sleep study in multiple sclerosis: correlations with clinical and brain magnetic resonance imaging findings. *J Neurological Sci* 1994; 125:194-7.
  59. Montplaisir J, Boucher S, Poirier G, Lavigne G, Lapierre O, Lesperance P. Clinical, polysomnographic, and genetic characteristics of restless legs syndrome: a study of 133 patients diagnosed with new standard criteria. *Movement Disorders* 1997; 12:61-5.
  60. Rosa RR, Bonnet MH. Reported chronic insomnia is independent of poor sleep as measured by electroencephalography. *Psychosomatic Med* 2000; 62:474-82.
  61. Zucconi M, Oldani A, Ferini-Strambi L, Smirne S. Arousal fluctuations in non-rapid eye movement parasomnias: the role of cyclic alternating pattern as a measure of sleep instability. *J Clin Neurophysiol* 1995; 12:147-54.

**Table 3A—Evidence Table for Studies of Patients with Insomnia**

AUTHOR	GRADE/ LEVEL	GROUPS	NUMBER	MALES	FEMALES	AGE	DX CRITERIA	Q1	Q2	Q3	RESULTS &/OR CONCLUSION
Vgontzas <sup>24</sup>	III/C	Insomnia (I) Control (C)	375 150	187 62	188 88	43 44	DSM3R ASDA 1979	N	na	N	1- SRBD and PLMD equally prevalent in I and C 2- REM changes had 21% PPV for depression
Zorick <sup>27</sup>	III/C	Insomnia (I) Normal Controls (C)	84 20	49 9	38 11	51 49	ASDA 1979	Y	na	Y	1- Dx categories within I could be determined by PSG 2- I: ↑ objective PSG evidence or DIMS vs. C
Rosa <sup>60</sup>	III/C	Insomnia (I) <sup>2</sup> Non-insomnia (C)	121 56	82 35	39 21	36 35	none	N	N	N	1- No PSG difference in I vs. C except for S2L 2- I: ↑ subjective SL, WASO, #wakes
Lichstein <sup>25</sup>	III/C	Media-recruited individuals with Primary Insomnia. (excluded likely SRBD, hypnotic Rx use, psychiatric disorder, PLMD, RLS)	80	32	48	69	ICSD DSM4	Y	na	Y	1- 23 patients had SRBD with AHI ≥ 15 (29%) 2- 34 patients had SRBD with AHI ≥ 5 (43%) 3- 3 patients had PLMD with PLMAI ≥ (4%) 4- There is a high incidence of occult sleep apnea in elderly who appear to have primary insomnia
Rochrs <sup>26</sup>	III/C	SRBD with insomnia (I) SRBD with sleepiness (S)	16 65	2 63	14 2	45 47	none	Y	na	Y	1- Patients with insomnia may have SRBD 2- I: ↓ AI, less ↓ SaO <sub>2</sub> , ↑ central episodes vs. S 3- Sex Differences: ↑ ρ with insomnia, ↑ σ with sleepiness
Dorsey <sup>57</sup>	IV/C	Objective insomnia (OI) Subjective insomnia (SI) Control (C)	9 9 13	? ? ?	? ? ?	? ? ?	SL>45 mins <sup>1</sup>	Y	na	N	1- No group differences in PSG measures or MSLT SL 2- SI did not estimate MSLT SL accurately 3- OI did estimate MSLT SL accurately 4- ↑ neuroticism for SI, ↑ introversion for OI
Reynolds <sup>28</sup>	IV/C	Elderly patients with complaints of insomnia and daytime sleepiness	27	18	9	63	ASDA 1979	Y	na	Y	1- DIMS patients have depression (33%) and psychophysiological insomnia 2- DOES patients were more likely to have SRBD 3- ↓REML in patients with depression
Edinger <sup>29</sup>	V/D	Patients with insomnia presenting at sleep clinic	100	46	54	46	ASDA 1979	Y	N	Y	1- PSG provided important Dx info on 65% of sample 2- 34% sample Dx's essentially by PSG (25% PLMD, 3% apnea, 6% subjective insom) 3- PSG ruled out PLMD in 9% and apnea in 7% 4- PSG useful in patients 40 years and older

**Notation-** Q1: Did polysomnography help in the evaluation of insomnia in this study? Q2: Did polysomnography provide information helpful for understanding treatment failure? Q3: Did polysomnography differentiate between insomnia of different etiologies? PSG- Polysomnography, PLMD- Periodic Limb Movement Disorder, SRBD- Sleep-related Breathing Disorder, SEI- Sleep Efficiency Index, SL- Sleep Latency, WASO- Wake After Sleep Onset, TST- Total Sleep Time, SWS- Slow Wave Sleep, S2- Stage 2 Sleep, S2L- Latency to stage 2 sleep, REML- Latency to REM Sleep, REMD- REM density, ICSD- International Classification of Sleep Disorders, DSM- Diagnostic and Statistical Manual, ASDA 1979- American Sleep Disorders 1979 Sleep-wake disorder nosology. MSLT - multiple sleep latency test, DIMS - disorders of initiation and maintenance of sleep, RLS - restless leg syndrome, PPV- positive predictive value, PLMAI - periodic limb movement arousal index (number of PLMA per hour of sleep), DOES - disorders of excessive somnolence.

<sup>1</sup> These were college students volunteering for credit, they were not a self-selected clinical group seeking treatment

<sup>2</sup> Patients with AI>10 excluded from sample

**Table 3B**—Evidence Table for Studies of Patients with Psychiatric Disorders

AUTHOR	GRADE / LEVEL	GROUPS	NUMBER	MALES	FEMALES	AGE	DX CRITERIA	Q1	Q2	Q3	RESULTS &/OR CONCLUSION
Thase <sup>40</sup>	III/C	Depressed Outpatients (O)	181	66	115	37	DSM3R	N	Y	N	1-O and I: ↑REMD, ↓REML, ↓SEI vs. C
		Depressed Inpatients (I)	44	19	25	33					2- REM markers readily distinguish depressed from C
		Controls (C)	44	16	28	32					3- Abnormal sleep associated with poor treatment response
Merica <sup>47</sup>	IV/C	Depression, Major (M)	48	?	?	47	DSM3R	Y	N	N	1- All 3 groups had similar sleep fragmentation
		Dysrhythmia (D)	36	?	?	43					2- REM period duration increase across night blunted in group with Major Depressive Disorder
		Insomnia (I)	27	?	?	37					
		Age-matched controls (C)	?	?	?	?					
Arriaga <sup>41</sup>	IV/C	Panic+Agoraphobia (P)	14	6	8	34	DSM4	na	na	na	1- P: ↓SEI, ↑Wake%, ↓S4%, ↓SWS vs. C
		Age & sex matched normal controls (C)	14	6	8	34					2- No group difference for TST, SL, S1%, S2%, S3%, REM%
Lauer <sup>46</sup>	IV/C	Panic Disorder (P)	22	8	14	33	DSM3R	Y	na	Y	1- P: ↓REML, ↑SL vs. C
		Depression, Major (D)	12	4	8	30					2- D: ↓SEI vs. C
		Age-matched controls (C)	12	4	8	31					3- P: ↑SL vs. D
Pecknold <sup>48</sup>	IV/C	Panic+agoraphobia (P)	43	18	26	33	DSM3	N	N	N	1- SEI: A=P, A<C, P<C
		Anxiety Disorder (A)	11	?	?	?					2- REM: no differences
		Historical controls (C)	57	?	?	?					3- S2: A=P, A<C, P<C 4- SWS: A=P, A>C, P>C 5- TST: A>P, A=C, P<C
Stein <sup>49</sup>	IV/C	Panic Disorder (P)	16	5	11	37	DSM3R	N	na	N	1- P: ↓Δ sleep, ↓TST vs. C
		Control (C)	16	?	?	37					2- C: ↑S2% vs. P
Hohagen <sup>44</sup>	IV/C	Obsessive Compulsive Disorder (OCD)	22	10	12	39	DSM3R	na	na	na	1- OCD: ↓SEI, ↑Wake% vs. C
		Age-matched controls (C)	22	12	10	39					2- No group difference in sleep macroarchitecture
Hurwitz <sup>45</sup>	IV/C	Combat Vets with PTSD	18	18	0	45	DSM3R	Y	na	na	1- ↑SL to S2, ↓arousals/hour from SWS on night2, ↓ subjective estimates of TST, ↑ sleep state misperception in PTSD vs. C
		Noncombat Vet controls (C)	10	10	0	47					2- No group difference in sleep macroarchitecture

**Notation**- Q1: Did polysomnography help in the evaluation of insomnia in this study? Q2: Did polysomnography provide information helpful for understanding treatment failure? Q3: Did polysomnography differentiate between insomnia of different etiologies? PSG- Polysomnography, PLMD- Periodic Limb Movement Disorder, SRBD- Sleep-related Breathing Disorder, SEI- Sleep Efficiency Index, SL- Sleep Latency, WASO- Wake After Sleep Onset, TST- Total Sleep Time, SWS- Slow Wave Sleep, S2- Stage 2 Sleep, S2L- Latency to stage 2 sleep, REML- Latency to REM Sleep, REMD- REM density, ICSD- International Classification of Sleep Disorders, DSM- Diagnostic and Statistical Manual, ASDA 1979- American Sleep Disorders 1979 Sleep-wake disorder nomenclature. MSLT - multiple sleep latency test, DIMS - disorders of initiation and maintenance of sleep, RLS - restless leg syndrome, PPV- positive predictive value, PLMAI - periodic limb movement arousal index ( number of PLMA per hour of sleep), DOES - disorders of excessive somnolence

**Table 3C—Evidence Table for Studies of Patients with Sleep Disorders, Neurological Disorders, and Other Disorders**

AUTHOR	GRADE / LEVEL	GROUPS	NUMBER	MALES	FEMALES	AGE	DX CRITERIA	Q1	Q2	Q3	RESULTS &/OR CONCLUSION
Zucconi <sup>61</sup>	III/C	Parasomnia DOA <sup>3</sup> (A) Parasomnia, Other (O) Control (C)	13 8 6	9 4 6	4 4 0	22 23 25	ICSD	Y	na	N	1- 60% parasomnia events were during a cyclic alternating pattern (CAP) episode 2- Sleep Macroarchitecture did not differ between groups
Allen <sup>50</sup>	IV/C	Demented (D) Non-demented (ND)	30 14	13 7	17 7	80 83	none	na	na	na	1- D: ↓ TST, ↓ S2, ↓ REM, ↑ arousals vs. ND 2- No sleep difference between D subtypes 3- No 1 <sup>st</sup> night effect for D
Ferini-Strambi <sup>58</sup>	IV/C	Multiple Sclerosis (MS) Controls (C)	25 25	13 13	12 12	40 40 <sup>1</sup>	DSM-III-R	na	na	na	1- MS: ↑ PLMD (n=9) vs. C (n=2) 2- MS: ↓ SEI, ↑ #wakes vs. C 3- No group difference in sleep macroarchitecture
Manu <sup>52</sup>	IV/C	Chronic Fatigue (CF) <sup>2</sup> Subgroup with αΔ sleep Subgroup without αΔ sleep	30 8 22	8 2 7	21 6 15	41 44 40	none	Y	na	Y	1- αΔ sleep identified in 8 patients with CF (26%) 2- αΔ sleep more common in CF without depression 3- 10 CF patients other sleep disorders- PLMD (n=6), SRBD (n=4), NS (n=1).
Montplaisir <sup>59</sup>	IV/C	Restless Leg Syndrome	133	63	70	51	ICSD	N	N	N	1- RLS: ↑ SL, ↑ sleep disruption 2- 63% had 1 <sup>o</sup> family member with RLS 3- Comorbid PLMD common
Evans <sup>51</sup>	V/D	Patients with head injury	138	?	?	27	none	N	Y	N	1- Some utility for predicting outcome suggested

**Notation-** Q1: Did polysomnography help in the evaluation of insomnia in this study? Q2: Did polysomnography provide information helpful for understanding treatment failure? Q3: Did polysomnography differentiate between insomnia of different etiologies? PSG- Polysomnography, PLMD- Periodic Limb Movement Disorder, SRBD- Sleep-related Breathing Disorder, SEI- Sleep Efficiency Index, SL- Sleep Latency, WASO- Wake After Sleep Onset, TST- Total Sleep Time, SWS- Slow Wave Sleep, S2- Stage 2 Sleep, S2L- Latency to stage 2 sleep, REML- Latency to REM Sleep, REMD- REM density, ICSD- International Classification of Sleep Disorders, DSM- Diagnostic and Statistical Manual, ASDA 1979- American Sleep Disorders 1979 Sleep-wake disorder nosology. MSLT - multiple sleep latency test, DIMS - disorders of initiation and maintenance of sleep, RLS - restless leg syndrome, PPV- positive predictive value, PLMAI - periodic limb movement arousal index ( number of PLMA per hour of sleep), DOES - disorders of excessive somnolence

<sup>1</sup> Controls were age matched but the group age was not explicitly stated

<sup>2</sup> Consecutive patients in a cohort population

<sup>3</sup> Disorders of Arousal, including Confused Arousals, Sleep Walking, and Sleep Terror